



A comprehensive approach to esthetics, health and function.

8801 N. Meridian Street, Suite 103
Indianapolis, Indiana 46260
317.574.0600 voice / 317.574.0606 fax

9865 E. 116th Street, Suite 100
Fishers, Indiana 46037
317.842.2273 voice / 317.842.7911 fax

info@indyimplants.com / www.indyimplants.com

Patient Self-Care Following Sinus Augmentation Surgery

ACTIVITY: After leaving our dental office today, we suggest you consider relaxing or at least limiting your activity as much as possible for the remainder of the day. Avoid strenuous activity or aerobic exercise for the next 2-3 days.

DISCOMFORT: Some discomfort may be present when the anesthesia wears off. You have been given a prescription for an anti-inflammatory (NSAID) analgesic. Please take this as prescribed. This drug will greatly decrease the possibility of post-surgical swelling and pain and has been shown to accelerate healing. After the first 3 days, 2 Tylenol, or 2 Advil taken as indicated on the bottle, should keep you comfortable. Should intense discomfort occur at any time after the surgery, please take the narcotic medication (if one has been prescribed for you), as directed, or telephone the doctor and he will phone in a narcotic prescription for you. You may take both the narcotic and NSAID together. Their analgesic effect will be synergistic.

INFECTION: Please continue taking your antibiotic as prescribed until all of the pills have been taken. If you notice that after a few days, pain or swelling are increasing or that you are experiencing an elevated temperature, please call the doctor.

ANTIHISTAMINE USE: Please continue taking your antihistamine, one tablet every 5-8 hours, or 2-3 times per day for at least the first 4-5 days, particularly if you are having any symptoms of "stuffiness" or tendency to sneeze. Antihistamines can make you drowsy. If this is the case then reduce the dosage, i.e. take them twice a day instead of three times a day. Continue to use the nasal spray, one puff every 5-8 hours for the first three days, or longer if symptoms of "stuffiness" and difficulty breathing through your nose are present.

SWELLING: It is normal to experience some facial swelling after surgery. To minimize this, apply an ice bag wrapped in a light towel or handkerchief to the outside of your face over the operated area. This should be left on your face for about 20 minutes, then removed for 20 minutes, or alternated from side-to-side, between operated areas, for 2-3 hours after surgery. The use of both ice and the NSAID analgesic as described above will reduce facial swelling.

BLEEDING: It is common to have slight bleeding for a few hours following surgery. Most of the blood you may see in your mouth is actually a little bit of blood mixed with a lot of saliva - blood is a very strong "dye" and a little bit of blood will color your saliva dramatically. If bleeding persists, apply a moistened tea bag to the surgical site with gentle pressure for 15-20 minutes. Let yourself rest for 15 minutes then repeat the application of a tea bag for a second time if you still have slight bleeding. If excessive bleeding continues, please call our office or the doctor after normal office hours. You may experience some blood discharge from your nose. If so, lie down, with your head slightly elevated and apply an ice pack over your midface-nasal area. If this bleeding does not stop within 30 minutes, telephone the doctor.

ORAL HYGIENE: Brushing and oral hygiene procedures should be done as usual in all untreated areas. In exposed, operated areas limit your oral hygiene to brushing using the sponge swabs provided. In areas covered by periodontal dressing, brush only the chewing surfaces of teeth. Avoid dental flossing in operated areas during the first week following surgery. Also avoid the use of water irrigation devices such as Water-piks for 1 month following surgery. The exposed operated areas should be gently swabbed with a Q-tip (cotton tipped applicator) or sponge swab saturated with Peridex (Chlorhexidine) as a last oral hygiene procedure at bedtime and again in the morning, after eating and drinking.

EATING AND DRINKING: Do not try to eat until all anesthesia (numbness) has worn off. High protein foods and liquids are desirable for 3-5 days following surgery. Semi-solid foods may be eaten as long as this may be done comfortably. . Eggs, custard, yogurt, pasta, steamed vegetables, casseroles, cooked cereals are some things that you might consider eating during the first few days following your surgery. Avoid spicy, salty, acidic, or very hot foods or liquids. Also, avoid nuts, chips or other crunchy or fibrous foods which may become caught between your teeth.

We want your treatment to provide as positive a healing experience as possible. If you are having any problems or have any questions not covered in these guidelines, please feel free to call us at any time.