



## Implant Dentistry and Periodontics

Michael D. Edwards, DDS, MSD

**A comprehensive approach to esthetics, health and function.**

8801 N. Meridian Street, Suite 103  
Indianapolis, Indiana 46260  
317.574.0600 voice / 317.574.0606 fax

9865 E. 116th Street, Suite 100  
Fishers, Indiana 46037  
317.842.2273 voice / 317.842.7911 fax

info@indyimplants.com / www.indyimplants.com

### Patient Self-Care Following Soft Tissue Graft Surgery

**ACTIVITY:** After leaving our dental office today, we suggest you consider relaxing or at least limiting your activity as much as possible for the remainder of the day. Avoid strenuous activity and all aerobic exercise for the next 3 days as this might cause the graft to separate from the underlying tissue due to an elevation in your blood pressure.

**DISCOMFORT:** Some discomfort may be present when the anesthesia wears off. You have been given a prescription for an anti-inflammatory (NSAID) analgesic. Please take this as prescribed. This drug will greatly decrease the possibility of post-surgical swelling and pain and has been shown to accelerate healing. After the first 3 days, Tylenol, or Advil taken as indicated on the bottle, should keep you comfortable. Should intense discomfort occur at any time after the surgery, please telephone the doctor and he will phone in a prescription for you.

**SWELLING:** It is normal for some swelling to occur after surgery, particularly in the lower jaw. To minimize swelling after surgery, apply an ice bag wrapped in a light tea towel or handkerchief to the outside of your face over the operated area. This should be left on your face for about 20 minutes, then removed for 20 minutes, or alternated from side-to-side, between operated areas, for 2-3 hours after surgery. The use of both ice and the NSAID analgesic as described above will reduce the amount and duration of facial swelling.

**BLEEDING:** It is common to have slight bleeding for a few hours following soft tissue graft surgery. Try to avoid repeatedly spitting or rinsing your mouth during the first 3-4 hours after surgery as this will encourage bleeding. If bleeding persists, remove blood by wiping with a Kleenex tissue to identify the location of the bleeding site and then apply a moistened tea bag to the area with firm but gentle pressure for 15-20 minutes. Let yourself rest for 15 minutes then repeat the application of a tea bag for a second time if you still have slight bleeding. If excessive bleeding continues, please call us. Remember, most of the blood you may see in your mouth is actually a little bit of blood mixed with a lot of saliva - blood is a very strong "dye" and a little bit of blood will color your saliva dramatically.

**ORAL HYGIENE:** Brushing and oral hygiene procedures should be done as usual in all untreated areas starting the night of surgery. In exposed, operated areas limit your oral hygiene to gentle brushing using the sponge swabs provided. In areas covered by periodontal dressing, brush only the chewing surfaces of teeth. Avoid dental flossing in operated areas during the first week following surgery and in areas covered with periodontal dressing. Also avoid the use of water irrigation devices such as Water-piks or electric toothbrushes in the operated areas for 2-3 weeks following surgery. The exposed operated areas should be gently swabbed with a Q-tip (cotton tipped applicator) or sponge swab saturated with Peridex (Chlorhexidine) as a last oral hygiene procedure at bedtime and again in the morning, after eating and drinking.

**EATING AND DRINKING:** Do not try to eat until all anesthesia (numbness) has worn off. High protein foods and liquids are desirable for 3-5 days following surgery. Semi-solid foods may be eaten as long as this may be done comfortably. . Eggs, custard, yogurt, pasta, steamed vegetables, casseroles, cooked cereals are some things that you might consider eating during the first few days following your surgery. Avoid spicy, salty, acidic, or very hot foods or liquids. Also, avoid hard foods: toast crust, nuts, chips or crunchy or fibrous foods such as raw carrots that may become caught between your teeth or traumatize exposed operated tissues.

We want your treatment to provide as positive a healing experience as possible. If you are having any problems or have any questions not covered in these guidelines, please feel free to call us at any time.