



A comprehensive approach to esthetics, health and function.

8801 N. Meridian Street, Suite 103
Indianapolis, Indiana 46260
317.574.0600 voice / 317.574.0606 fax

9865 E. 116th Street, Suite 100
Fishers, Indiana 46037
317.842.2273 voice / 317.842.7911 fax

info@indyimplants.com / www.indyimplants.com

Patient Self-Care Following Gingival Treatment

ACTIVITY: After leaving our dental office, we suggest you consider relaxing or at least limiting your activity as much as possible for the remainder of the day. Avoid strenuous activity and aerobic exercise for the next 3 days.

DISCOMFORT: Some discomfort may be present when the anesthesia wears off. You have been given a prescription for an anti-inflammatory (NSAID) analgesic. Please take this as indicated on the bottle. This drug will greatly decrease the possibility of post-treatment swelling and pain and has been shown to accelerate healing. After the first 3 days, Tylenol, or Advil taken as indicated on the bottle, should keep you comfortable. Should intense discomfort occur at any time after the treatment, please take the narcotic medication (if one has been prescribed for you), as directed, or telephone the doctor. You may take both the narcotic and NSAID together. Their analgesic effect will be synergistic.

INFECTION: If you have been given an antibiotic please continue taking it, as directed, until all of the pills have been taken. If you notice that after a few days, pain or swelling are increasing or that you are experiencing an elevated temperature, please call the doctor.

SWELLING: It is normal for some swelling to occur after treatment, particularly in the lower jaw. To minimize swelling after treatment, apply an ice bag wrapped in a light towel or handkerchief to the outside of your face over the treated area. This should be left on your face for about 20 minutes, then removed for 20 minutes, or alternated from side-to-side, between treated areas, for 6 hours after treatment or until bedtime. The use of both ice and the NSAID analgesic as described above will reduce the amount and duration of facial swelling.

BLEEDING: It is common to have slight bleeding for a few hours following periodontal treatment. If bleeding persists, apply a moistened tea bag to the treatment site with firm but gentle pressure for 15-20 minutes. Let yourself rest for 15 minutes then repeat the application of a tea bag for a second time if you still have slight bleeding. If excessive bleeding continues, please call our office. Remember, most of the blood you may see in your mouth is actually a little bit of blood mixed with a lot of saliva - blood is a very strong "dye" and a little bit of blood will color your saliva dramatically.

ORAL HYGIENE: Brushing and oral hygiene procedures should be done as usual in all untreated areas. In exposed, treated areas limit your oral hygiene to brushing with the sponge swabs provided. In areas covered by periodontal dressing, brush only the chewing surfaces of teeth. Avoid dental flossing in treated areas during the first week following treatment. Avoid the use of water irrigation devices such as Water-piks for 2 weeks following treatment. Any exposed treated areas should be gently swabbed with a Q-tip (cotton tipped applicator) or sponge swab saturated with Peridex (Chlorhexidine) as a last oral hygiene procedure at bedtime and again in the morning, after eating and drinking.

EATING AND DRINKING: Limit diet to cool and soft foods until all anesthesia (numbness) has worn off. Semi-solid foods may be eaten as long as this may be done comfortably. Eggs, custard, yogurt, pasta, steamed vegetables, casseroles and cooked cereals are some things that you might consider eating during the first few days following your treatment. Avoid spicy, salty, acidic, or very hot foods or liquids. Also, avoid nuts, chips or other crunchy or fibrous foods that may become caught between your teeth.

CARE OF THE DRESSING: If small pieces of the dressing break off, do not be concerned. If large portions of the dressing are displaced or lost, and discomfort is experienced, please contact our office. If the dressing should fall off after 4-5 days, and you are comfortable, replacement of the dressing is unnecessary.

We want your treatment to provide as positive a healing experience as possible. If you are having any problems or have any questions not covered in these guidelines, please feel free to call us at any time.