

Patient Self-Care Following Implant Placement Treatment

ACTIVITY: After leaving our dental office, we suggest you consider relaxing or at least limiting your activity as much as possible for the remainder of the day. Avoid strenuous activity or aerobic exercise for the next 2-3 days.

DISCOMFORT: Some discomfort may be present when the anesthesia wears off. You have been given a prescription for an anti-inflammatory (NSAID) analgesic. Please take this as directed on the bottle. This drug will greatly decrease the possibility of post-treatment swelling and pain and has been shown to accelerate healing. After the first 3 days, Tylenol, or Advil taken as indicated on the bottle, should keep you comfortable. Should intense discomfort occur at any time after the treatment, please take the narcotic medication (if one has been prescribed for you), as directed, or telephone the doctor. You may take both the narcotic and NSAID together. Their analgesic effect will be synergistic.

INFECTION: If you have been given an antibiotic please continue taking it as prescribed until all of the pills have been taken. If you notice that after a few days, pain or swelling are increasing or that you are experiencing an elevated temperature, please call the doctor.

SWELLING: It is normal for some swelling to occur after treatment, particularly in the lower jaw. To minimize swelling after surgery, apply an ice bag wrapped in a light towel or handkerchief to the outside of your face over the treated area. This should be left on your face for about 20 minutes, then removed for 20 minutes, or alternated from side-to-side, between treated areas, for 2-3 hours after treatment. The use of both ice and the NSAID analgesic as described above will reduce the amount and duration of facial swelling.

BLEEDING: It is common to have slight bleeding for a few hours following treatment. If bleeding persists, apply a moistened tea bag to the treated site with firm but gentle pressure for 15-20 minutes. Let yourself rest for 15 minutes then repeat the application of a tea bag for a second time if you still have slight bleeding. If excessive bleeding continues, please call our office. Remember, most of the blood you may see in your mouth is actually a little bit of blood mixed with a lot of saliva - blood is a very strong "dye" and a little bit of blood will color your saliva dramatically.

ORAL HYGIENE: Brushing and oral hygiene procedures should be done as usual in all untreated areas. In exposed, treated areas limit your oral hygiene to brushing using the sponge swabs provided. In areas covered by periodontal dressing, brush only the chewing surfaces of teeth. Avoid dental flossing in treated areas during the first week following surgery. Also avoid the use of water irrigation devices such as Water-piks for 1 month following treatment. The exposed treated areas should be gently swabbed with a Q-tip (cotton tipped applicator) or sponge swab saturated with Peridex (Chlorhexidine) as a last oral hygiene procedure at bedtime and again in the morning, after eating and drinking.

EATING AND DRINKING: Limit diet to cool and soft foods until all anesthesia (numbness) has worn off. High protein foods and liquids are desirable for 3-5 days following treatment. Semi-solid foods may be eaten as long as this may be done comfortably. Eggs, custard, yogurt, pasta, steamed vegetables, casseroles and cooked cereals are some things that you might consider eating during the first few days following your surgery. Avoid spicy, salty, acidic, or very hot foods or liquids. Also, avoid nuts, chips or other crunchy or fibrous foods which may become caught between your teeth.

USE OF REMOVABLE DENTAL APPLIANCES: If you normally wear a removable appliance which replaces missing teeth, and it rests on the treated area where implants have been placed, do not use this appliance as any pressure on the treatment site could be detrimental to healing of the implants as well as cause discomfort.

CARE OF THE DENTAL APPLIANCE: It should be brushed and rinsed when you perform your normal oral hygiene care. If a soft "cushioning" liner has been applied, it is essential that this be checked frequently to see if it is hardening and needs replacement. If it hardens and rests on the implant treatment site, it could be detrimental to healing of the implants.

We want your treatment to provide as positive a healing experience as possible. If you are having any problems or have any questions not covered in these guidelines, please feel free to call us at any time.